INTERNATIONAL JOURNAL OF RESEARCH IN SCIENCE & TECHNOLOGY

e-ISSN:2249-0604; p-ISSN: 2454-180X

Identification, Isolation and Confirmation of Alkaloids in Rare Medicinal Plants of Telangana

Dr. K. Venkata Krishna

Assistant Professor of Chemistry

Government Degree College for Women (A), Nalgonda, Telangana, India

Paper Received: 30th July 2021; **Paper Accepted:** 05th October 2021;
Paper Published: 09th November 2021

DOI: http://doi.org/10.37648/ijrst.v11i04.006



How to cite the article:

Dr K. Venkata Krishna, Identification, Isolation and Confirmation of Alkaloids in Rare Medicinal Plants of Telangana, IJRST, Oct-Dec 2021, Vol 11, Issue 4, 52-61, DOI: http://doi.org/10.37648/ijrst.v11i04.006

ABSTRACT

Alkaloids are nitrogen-containing compounds found in many medicinal plants. Several rare or less-studied species in Telangana contain these compounds and are used by local communities. This paper describes how workers identify, isolate, and confirm alkaloids from such plants. It covers extraction steps, acid-base partitioning, chromatography, thin-layer and high-performance liquid chromatography, and spectroscopic methods. Examples include vasicine from Adhatoda, solasodine from Solanum species, and other bases reported from the region. Tables list plants, plant parts, and typical alkaloid yields. Chemical structures and reaction equations appear where they help explain the tests. The methods allow researchers to link traditional use with measurable chemical content when plant material is collected carefully and identified correctly.

Keywords: *alkaloids; medicinal plants; Telangana; isolation; identification; TLC; HPLC; spectroscopy*

I. INTRODUCTION

Telangana lies on the Deccan Plateau and supports many plant species used in local medicine. Ethnobotanical surveys in Nalgonda, Warangal, and nearby districts list hundreds of taxa employed for common ailments. Among these, a smaller number contain alkaloids that account for much of the observed activity. Alkaloids are basic compounds that form salts with acids. This property allows simple separation from neutral and acidic plant constituents.

Rare or less-collected species still grow in forest patches and along field margins. Local healers continue to use them, yet published chemical data remain limited. Systematic isolation and confirmation help verify traditional claims and guide conservation. The present paper outlines practical laboratory steps that have been applied to Indian medicinal plants and that can be adapted to material collected in Telangana.

II. SELECTED PLANTS AND THEIR ALKALOIDS

Adhatoda vasica (syn. Justicia adhatoda) grows widely in Telangana and is used for respiratory complaints. Its leaves contain the quinazoline alkaloid vasicine (peganine) together with vasicinone. Solanum species such as S. xanthocarpum and S. nigrum yield steroidal alkaloids including solasodine and its glycosides. Catharanthus roseus, cultivated and naturalized in the region, produces indole alkaloids such as vincristine and vinblastine. Rauvolfia

serpentina and related species supply reserpine-type bases. These plants illustrate the range of alkaloid classes present in the local flora.




Figure 1. Structure of vasicine (peganine), a quinazoline alkaloid from Adhatoda vasica.




Figure 2. Structure of solasodine, a steroidal alkaloid from Solanum species.




Figure 3. Structure of berberine, a protoberberine alkaloid found in several Indian medicinal plants.

A. ***Table 1. Examples of alkaloid-containing plants reported from Telangana and surrounding areas***

Plant	Main alkaloid class	Plant part used	Local use note
Adhatoda vasica	Quinazoline	Leaves	Respiratory
Solanum xanthocarpum	Steroidal	Whole plant, fruit	Cough, fever
Catharanthus roseus	Indole	Leaves, root	Various

III. COLLECTION AND EXTRACTION

Workers collect plant material during the flowering or fruiting season when alkaloid levels are often highest. They dry the material in shade to limit degradation and grind it to a coarse powder. Extraction usually employs methanol or ethanol at room temperature or under gentle reflux. The combined extract is filtered and concentrated under reduced pressure.

Acid-base partitioning concentrates the alkaloids. The concentrated extract is suspended in dilute acid (commonly 1–2 % acetic or hydrochloric acid). Neutral and acidic compounds remain in the organic layer or precipitate. The aqueous acid solution is basified with ammonia or sodium carbonate to pH 9–10. The free bases are then extracted into chloroform or dichloromethane. Evaporation of the organic solvent yields the crude alkaloid fraction.




Figure 4. Schematic workflow for isolation of alkaloids from plant material.

The overall recovery can be expressed simply as:

Crude alkaloid yield (%) = (mass of crude fraction / mass of dry plant material) × 100

IV. Chromatographic Separation

Column chromatography on silica gel or alumina separates the crude mixture. Workers elute with gradients of chloroform-methanol or hexane-ethyl acetate. Fractions are monitored by thin-layer chromatography (TLC). Dragendorff's reagent or Mayer's reagent visualizes the alkaloid spots as orange or cream precipitates.

Preparative TLC or repeated column runs yield pure compounds. High-performance liquid chromatography (HPLC) on reverse-phase columns with ultraviolet detection provides both analytical and preparative separation. Retention times are compared with authentic standards when available.

V. Confirmation of Identity

Chemical tests give a first indication. Dragendorff's reagent (potassium bismuth iodide) produces an orange-red precipitate with most alkaloids. Mayer's reagent (mercuric potassium iodide) gives a cream precipitate. Wagner's reagent (iodine in potassium iodide) yields a brown precipitate. These tests are not specific but confirm the presence of basic nitrogen compounds.

Spectroscopic methods provide definitive identification. Ultraviolet spectra show characteristic absorption maxima. Infrared spectra reveal functional groups such as

carbonyl or hydroxyl. Proton and carbon-13 nuclear magnetic resonance spectra establish the carbon skeleton and relative configuration. Mass spectrometry supplies the molecular ion and fragmentation pattern. Comparison with literature data or co-injection with a standard confirms the structure.

For vasicine the molecular formula is C11H12N2O. The mass spectrum typically shows a molecular ion at m/z 188. Characteristic fragment ions arise from loss of the side-chain hydroxyl group. Solasodine (C27H43NO2) gives a molecular ion near m/z 413 and fragments typical of steroidal alkaloids.

VI. Practical Considerations for Telangana Material

Local plants may contain lower or more variable alkaloid levels than cultivated material from other regions. Soil, rainfall, and collection season affect content. Workers therefore analyze several batches and report ranges rather than single values. Proper botanical identification by a taxonomist prevents confusion with similar species that lack the target alkaloids.

Conservation status must be checked before large collections. Some species listed in ethnobotanical surveys are uncommon outside protected forest areas. Sustainable harvesting or cultivation trials can supply material for further study without depleting wild populations.

Workers also examine the distribution of alkaloids within the plant. Leaves often contain higher levels of vasicine in Adhatoda, while roots of Rauvolfia accumulate reserpine-type compounds. Fruits of Solanum species are rich in solasodine glycosides. Sampling different organs helps locate the best starting material and avoids unnecessary collection of low-yielding parts.

Drying conditions affect recovery. Shade drying at ambient temperature preserves heat-sensitive bases better than sun drying. Over-drying or high humidity can promote enzymatic degradation or mold growth that reduces alkaloid content. Workers therefore monitor moisture and store dried material in airtight containers away from light.

Solvent choice influences both yield and selectivity. Methanol extracts a broader range of compounds than chloroform. Ethanol is preferred when residual solvent toxicity is a concern. Sequential extraction with solvents of increasing polarity can fractionate the extract before acid-base partitioning and reduce the load on later chromatographic steps.

pH control during basification is critical. Incomplete basification leaves alkaloids in the aqueous phase. Excessive base can cause emulsion formation or hydrolysis of sensitive glycosides. Most workers adjust to pH 9–10 with dilute ammonia and extract immediately to minimize degradation.

Column chromatography parameters affect resolution. Particle size of silica gel, column diameter, and flow rate determine separation quality. Gradient elution from non-polar to polar solvents separates compounds of different polarity. Fractions of 10–20 mL are collected and combined only after TLC confirms identical composition.

HPLC method development begins with a C18 column and a methanol-water or acetonitrile-water gradient containing a small amount of acid or buffer to improve peak shape for basic compounds. Detection at 254 or 280 nm suits most aromatic alkaloids. Quantification requires calibration curves prepared from pure standards.

Mass spectrometry provides molecular weight and structural clues. Electron-impact ionization gives extensive fragmentation useful for comparison with library spectra. Soft ionization methods such as electrospray produce prominent molecular ions that confirm molecular formula when high-resolution data are available.

Nuclear magnetic resonance spectra resolve stereochemistry. The chemical shifts and coupling constants of protons adjacent to the nitrogen atom or oxygen functions distinguish isomers. Two-dimensional experiments (COSY, HSQC, HMBC) establish connectivity when one-dimensional spectra are crowded.

Comparison with authentic samples remains the most direct confirmation. When standards are unavailable, literature melting points, optical rotations, and spectral data serve as references. Discrepancies prompt re-examination of purity or consideration of previously unreported compounds.

Seasonal variation is large in many species. Alkaloid content in Adhatoda leaves peaks during the monsoon and declines in the dry season. Collectors therefore record collection date, location, and phenological stage so that later workers can reproduce or explain differences in yield.

Ethnobotanical context guides selection of species. Surveys in Nalgonda and Warangal document dozens of plants used for cough, fever, and skin conditions. Cross-checking these lists against known alkaloid-bearing genera narrows the candidates for

laboratory study and increases the chance of finding active compounds.

Safety precautions are essential. Many alkaloids are biologically active at low doses. Workers handle extracts and pure compounds in a fume hood, wear gloves, and avoid skin contact or inhalation of dust. Waste solutions are neutralized and disposed of according to institutional guidelines.

Documentation includes voucher specimens deposited in a recognized herbarium. Without vouchers, later identification of the source plant becomes impossible if taxonomic revisions occur. Photographs of the living plant and notes on habitat further support reproducibility.

Biosynthetic pathways explain the structural diversity observed in local plants. Quinazoline alkaloids such as vasicine arise from anthranilic acid and ornithine-derived units. Steroidal alkaloids of the Solanum type form from cholesterol through a series of oxidations and nitrogen incorporation. Indole alkaloids in Catharanthus originate from tryptamine and secologanin. Understanding these routes helps predict which related species may contain similar compounds and guides targeted extraction.

Quantitative estimation often employs spectrophotometric methods after color development with reagents such as Dragendorff's or bromocresol green. Calibration curves prepared from pure vasicine or solasodine allow calculation of total alkaloid content in crude extracts. HPLC with external standards provides more specific quantification of individual compounds and detects co-occurring bases that TLC may miss.

Purity assessment combines melting-point determination, thin-layer chromatography in at least two solvent systems, and spectroscopic homogeneity. A single spot on TLC and a sharp melting point indicate high purity. Impurities appear as additional spots or broadened NMR signals. Recrystallization from methanol, ethanol, or acetone–water mixtures improves purity before final spectroscopic confirmation.

Storage of isolated alkaloids requires protection from light, moisture, and air. Many bases oxidize or form salts on standing. Workers keep pure compounds in amber vials under nitrogen or in a desiccator. Solutions for analysis are prepared fresh to avoid degradation that alters chromatographic retention times or spectral features.

Comparative studies of material from different districts of Telangana reveal geographic variation. Plants collected near Nalgonda may differ in alkaloid profile

from those in Warangal or Adilabad because of soil type, altitude, and associated vegetation. Recording GPS coordinates and soil characteristics with each collection allows later correlation of chemical data with environmental factors.

Traditional preparation methods provide clues for laboratory extraction. Decoctions and infusions used by local healers extract water-soluble alkaloid salts. Alcoholic extracts capture free bases. Laboratory acid-base partitioning therefore mirrors, in a controlled way, the selective extraction practiced in ethnomedicine and helps explain why certain preparations show activity.

Safety and regulatory aspects deserve attention when alkaloids are isolated for further study. Some compounds possess potent pharmacological or toxic effects. Researchers follow institutional biosafety guidelines, label all fractions clearly, and restrict access to pure samples. Documentation of yields and spectral data supports any later patent or publication claims.

Future work can expand the survey to additional rare species listed in local floras but not yet examined chemically. Integration of field ethnobotany with laboratory isolation, followed by pharmacological screening of pure compounds, will strengthen the evidence base for traditional uses in Telangana and identify candidates for conservation or cultivation.

The overall process—from careful collection through extraction, separation, and spectroscopic confirmation—produces reliable chemical data. When these data are published with full experimental details and voucher information, other researchers can reproduce the findings and build upon them. This cumulative approach advances both phytochemistry and the sustainable use of medicinal plants in the region.

VII. CONCLUSION

Identification, isolation, and confirmation of alkaloids from rare medicinal plants of Telangana follow standard phytochemical procedures. Acid-base extraction, chromatography, and spectroscopic comparison allow workers to obtain pure compounds and match them to known structures. When these steps are combined with careful plant identification and seasonal collection, the chemical data support traditional uses and help prioritize species for further pharmacological or conservation work.

REFERENCES

1. Sreeramulu, N., Suthari, S., Ragan, A., & Raju, V. S. (2013). Ethno-botanico-

medicine for common human ailments in Nalgonda and Warangal districts of Telangana, Andhra Pradesh, India. Annals of Plant Sciences, 2(8), 1–10.

2. Bhakuni, D. S. (1984). Alkaloids from Indian medicinal plants and their biosynthesis. Proceedings of the Indian Academy of Sciences – Chemical Sciences, 93(4), 661–676.
3. Cordell, G. A. (1981). Introduction to alkaloids: A biogenetic approach. Wiley-Interscience.
4. Harborne, J. B. (1998). Phytochemical methods: A guide to modern techniques of plant analysis (3rd ed.). Chapman & Hall.
5. Wagner, H., & Bladt, S. (1996). Plant drug analysis: A thin layer chromatography atlas (2nd ed.). Springer.
6. Suthari, S., & Raju, V. S. (2018). Ethnobotanical explorations in Telangana, the youngest state in Union of India: A synoptic account. In M. Ozturk & K. R. Hakeem (Eds.), Plant and human health (Vol. 1, pp. 65–123). Springer.
7. Bruneton, J. (1999). Pharmacognosy, phytochemistry, medicinal plants (2nd ed.). Intercept.
8. Dewick, P. M. (2009). Medicinal natural products: A biosynthetic approach (3rd ed.). Wiley.
9. Roberts, M. F., & Wink, M. (Eds.). (1998). Alkaloids: Biochemistry, ecology, and medicinal applications. Plenum Press.
10. Mann, J. (1987). Secondary metabolism (2nd ed.). Oxford University Press.